

# Multiplicity and membranes collision in modified AdS

Pozdeeva E.O.

SINP MSU


based on I.Ya. Aref'eva, E.O. Pozdeeva, T.O.Pozdeeva

*Theor.Math.Phys. 176(1)(2013)*

II Russian-Spanish Congress

Particle and Nuclear Physics at all Scales and Cosmology


Saint-Petersburg, October 1-4, 2013



# QGP

Quark Gluon Plasma (QGP) has been discovered in Au+Au collision at energy 100 GeV for nucleon in 2005 @ RHIC

## QGP formation




# 5D gravity and 4D field theory are related

In holographic approach classical gravity in  $AdS_5$  describes strong coupling field theory in 4D Minkowski space

There is hypothesis that QGP formation in 4D space corresponds to Black Holes creation in dual 5D space.

Gubser, Klebanov, Polyakov, 9802109

Witten, 9802150




The gravitational shock wave in  $AdS_5$  space is dual to ultrarelativistic heavy-ion in 4D space-time.

Thus,


- heavy-ion collisions can be represented such as gravitational shock waves collisions in  $AdS_5$
- QGP formation is equivalent BH creation in  $AdS_5$

Gubser et al.; 0805.1551, 0902.4062



## Problem:

- How to get experimental dependence of multiplicity on energy from holographic model.
- Simplest holographic model is related with use of  $\mathcal{N} = 4$  SYM  
[But QCD is not SYM]
- Our goal: to study more complicate models to fit experimental data.



# Multiplicity and trapped surface area


Main conjecture: multiplicity is proportional to entropy

$$S \sim N$$

Gubser et al.; 0805.1551

On experiments can be measured only  $N_{ch}$  :  $N \sim N_{ch}$

B. B. Back et. al., 0210015[nucl-ex].



Accordingly experiment the charged-particles pseudorapidity density depends on colliding energy

$$dN_{ch}/d\eta \propto s_{NN}^{0.15}$$

for Pb-Pb and Au-Au collisions

$$dN_{ch}/d\eta \propto s_{NN}^{0.11} \quad \text{pp collision}$$


$E = (1/2)\sqrt{s_{NN}}$  - colliding energy for nucleon

K. Aamodt et al. [ALICE Collaboration], 1011.3916 [nucl-ex].

DISCREPANCY

The simple holographic model gives

$$dN_{ch}/d\eta \propto s_{NN}^{2/3}$$




The minimal black hole entropy can be estimated by trapped surface area

$$S \geq S_{trapped} = A_{trapped} / 4G_N$$

The trapped surface is surface whose null normals all propagate inward.

S. W. Hawking and D. Page, Thermodynamics Of Black Holes In Anti-de Sitter Space, Commun. Math. Phys. 87 (1983) 577.

C. S. Pe,ca, J. P. S. Lemos, 9805004 [gr-qc]






- $\mathcal{N}=4$  SYM is not QCD
- For holographic description of QCD a modified  $AdS_5$  is used to study the dependence of entropy on energy

Gursoya, Kiritsis et al., 0707.1324, 0707.1349

- Early the modification of  $AdS_5$  space-time by introduction of wrapping factor was applied for shock waves

Kiritsis, 1111.1931




- The collisions of shock waves with masses averaged over transversal surfaces are named wall-wall collisions.

- We describe heavy-ion collisions by the wall-wall shock wave collisions in  $AdS_5$  (or in its modification)

S. Lin, E. Shuryak, 1011.1918[hep-th]

I. Y. Arefeva, A. A. Bagrov and E. O. Pozdeeva,  
Holographic phase diagram of quark-gluon plasma  
formed in heavy-ions collisions," JHEP 1205, 117 (2012)

- In the modified 5D spaces we consider the wall-wall collisions.



# Einstein equation

The Einstein equation for particle in dilaton field has the form:

$$\left( R_{\mu\nu} - \frac{g_{\mu\nu}}{2} R \right) - \frac{g_{\mu\nu}}{2} \left( -\frac{4}{3} (\partial\Phi_s)^2 + V(\Phi_s) \right) - \frac{4}{3} \partial_\mu \Phi_s \partial_\nu \Phi_s - g_{\mu\nu} \frac{d(d-1)}{2L^2} = 8\pi G_5 J_{\mu\nu},$$


where  $J_{++} = \frac{E}{b^3(z)} \delta(x^1) \delta(x^2) \delta(z - z_*) \delta(x^+).$

Shock wave metric modified by wrapping factor

$$ds^2 = b^2(z) (dz^2 + dx^i dx^i - dx^+ dx^- + \phi(z, x^1, x^2) \delta(x^+) (dx^+)^2)$$

Aref'eva I.Ya. 0912.5482[hep-th]


M. Hotta, M. Tanaka, Shock-wave geometry with nonvanishing cosmological constant, *Class. Quantum Grav.* **10**, 307, 1993



Using shock ansatz we reduce the Einstein equation to the differential equation for shock wave profile and two equations defining the connection of field and field potential with b-factor:

$$\left( \partial_{x^1}^2 + \partial_{x^2}^2 + \partial_z^2 + \frac{3b'}{b} \partial_z \right) \phi(z, x_\perp) = -16\pi G_5 \frac{E}{b^3} \delta(x^1) \delta(x^2) \delta(z_* - z)$$

$$V(\Phi_s) = \frac{3}{b^2} \left( \frac{b''}{b} + \frac{2(b')^2}{b^2} - \frac{4b^2}{L^2} \right) \quad \Phi'_s = \pm \frac{3}{2} \sqrt{\left( \frac{2(b')^2}{b^2} - \frac{b''}{b} \right)}$$



# Shock wave profile equation for flat objects with masses uniformly distributed collisions

- mass distributed over the domain wall


$$\left( \partial_z^2 + 3 \frac{b'}{b} \partial_z \right) \phi^W(z) = -16\pi G_5 \frac{E}{b^3} \delta(z_* - z)$$

- mass distributed over the finite region with radius L

$$\left( \partial_z^2 + \frac{3b'}{b} \partial_z \right) \phi^w(z) = -16\pi G_5 \frac{E^*}{b^3} \delta(z_* - z), \quad E^* = \frac{E}{L^2}$$

- The equations coincide up to a constant factor  $L^2$
- The solutions to equations coincide up to constant too  $L^2$

$$\phi^w(z) = \frac{\phi^W(z)}{L^2}$$




# Trapped surface

- We apply the profile of shock wave with averaged mass for consideration of the black hole formation in AdS<sub>5</sub>.
- We identify the black hole creation with trapped surface formation.
- In the case with averaged mass there is condition to the shock wave profile at the trapped surface (in boundary points)

$$(\partial_z \phi^w)^2 |_{TS} = 4$$

- The trapped surface formation condition we will apply to obtain trapped surface boundary points




The trapped surface area is calculated as follows

$$S_{trap} = \frac{1}{2G_5} \int_C \sqrt{\det|g_{AdS_3}|} dz d^2x_{\perp}$$

where  $\det|g_{AdS_3}|$  is the metric determinant of  $AdS_3$

The relative area  $s$  depend with the formula

$$s = \frac{S_{trap}}{\int d^2x_{\perp}}$$



# AdS space-time modification

- For the standard AdS space-time  $b(z)$  factor has form  $b(z)=L/z$
- We modify AdS space-time using wrapping factors types


$$b = \left( \frac{L}{z - z_0} \right)^a$$

$$b = \frac{L}{z} \exp \left( -\frac{z^2}{R^2} \right)$$

$$b = \exp \left( -\frac{z}{R} \right)$$

$$b = \left( \frac{L}{z} \right)^a \exp \left( -\frac{z^2}{R^2} \right)$$

where  $a > 0$ ,  $R = 1$  fm,  $L \approx 4.4$  fm






# Relations between trapped surface boundary and collision points


Using the solution to general form of domain equation (for any wrapping factor) and the trapped surface conditions we obtain the relations

$$F(z_*) = \frac{b^{-3}(z_b)F(z_a) + b^{-3}(z_a)F(z_b)}{b^{-3}(z_a) + b^{-3}(z_b)} \quad ; \partial_z F(z) = b^{-3}(z)$$



$$b^{-3}(z_a) = \frac{b^{-3}(z_b)}{\frac{8\pi G_5 E}{L^2} b^{-3}(z_b) - 1}$$

between trapped surface boundary ( $z_a < z_b$ ) points and collision point  $z_*$



# Exponential wrapping factor


For the wrapping factor  $b = \exp\left(-\frac{z}{R}\right)$  we have obtained

following relations between boundary points and collision point

$$Z_A = \frac{L^2}{16\pi G_5 E} \cdot \frac{Z_B}{Z_B - \frac{L^2}{16\pi G_5 E}}, \quad Z_0 = \frac{L^2}{8\pi G_5 E}$$

$$Z_0 = \exp\left(\frac{3z_*}{R}\right), \quad Z_A = \exp\left(\frac{3z_a}{R}\right), \quad Z_B = \exp\left(\frac{3z_b}{R}\right)$$

For the considered case the collision point is fixed by energy.




The relative area of trapped surface dened by

$$s = \frac{3}{2RG_5} \left( \frac{1}{\exp\left(\frac{3z_a}{R}\right)} - \frac{1}{\exp\left(\frac{3z_b}{R}\right)} \right) = \frac{3}{2RG_5} \left( \frac{1}{Z_A} - \frac{1}{Z_B} \right)$$

The maximum entropy value is obtained for  $Z_b \gg 1$ , in this approximation

$$Z_a \sim \frac{L^2}{16\pi G_5 E}, \quad s \sim \frac{24\pi E}{RL^2}$$

The entropy dependence on energy is linear for the exponential wrapping factor



# Power-law wrapping factor


Power-law b-factor  $b = (L/z)^a$  gives following low boundary point and the collision point

$$z_A = \left( \frac{z_B^{3a}}{-1 + z_B^{3a} C^2} \right)^{\frac{1}{3a}}, \quad z_* = \left( \frac{z_A^{3a} z_B^{3a} (z_B + z_A)}{z_A^{3a} + z_B^{3a}} \right)^{\frac{1}{3a+1}}, \quad C^2 = \frac{8\pi G_5 E}{L^{3a+2}}$$

$$s = \frac{1}{2G_5(3a-1)} \left( z_A \left( \frac{L}{z_A} \right)^{3a} - z_B \left( \frac{L}{z_B} \right)^{3a} \right)$$

and relative area of trapped surface


The maximal entropy value will  
at  $Z_b \gg 1$  in assumption  $3a > 1$




The maximal entropy value is relate only with energy and constant parameters

$$s \Big|_{z_b \rightarrow \infty} = \frac{L^{3a}}{2G_5(3a-1)} z_A^{1-3a} = \frac{L}{2G_5} \left( \frac{8\pi G_5}{L^2} \right)^{\frac{3a-1}{3a}} E^{\frac{3a-1}{3a}}$$

For the power wrapping factor the entropy increase as  $E^{1-1/3a}$



The multiplicity of particles produced in collisions of heavy ions (PbPb-and AuAu-collisions) depends on energy as  $s_{NN}^{0,15} (E^{0.3})$  in the range  $10-10^3$  GeV.



K. Aamodt et al.  
 [ALICE Collaboration],  
 arXiv:1011.3916 [nucl-ex].

The model with power-law wrapping factor can coincide with experimental data at  $a \approx 0.47$

# Simple mixed factor

Mixed factor of the form  $b = \frac{L}{z} \exp\left(-\frac{z^2}{R^2}\right)$  gives the another relative area of trapped surface energy dependence

$$s = \frac{L^3}{2G_5} \left( -\frac{1}{2 \exp\left(\frac{3z_b^2}{R^2}\right) z_b^2} + \frac{1}{2 \exp\left(\frac{3z_a^2}{R^2}\right) z_a^2} + \frac{3 \text{Ei}\left(1, \frac{3z_b^2}{R^2}\right)}{2R^2} - \frac{3 \text{Ei}\left(1, \frac{3z_a^2}{R^2}\right)}{2R^2} \right)$$

which has the maximal value at  $z_b \rightarrow \infty$

$$s \Big|_{z_b \rightarrow \infty} = \frac{3}{4} \frac{L^3}{G_5 R^2} \left( -\text{Ei}\left(1, \frac{3z_a^2}{R^2}\right) + \frac{1}{3} \frac{R^2}{\exp\left(\frac{3z_a^2}{R^2}\right) z_a^2} \right)$$

and roughly is  $E^{\frac{2}{3}}(1 + 0.007 \ln E) - 3$  at  $10\text{GeV} \leq E < 1\text{TeV}$

# Complicate mixed factor

The wrapping factor


$$b = \left(\frac{L}{z}\right)^a \exp\left(-\frac{z^2}{R^2}\right)$$

gives the most complicate relative area of trapped surface energy dependence

$$s = \frac{F(z_B) - F(z_A)}{2G_5}$$

$$F(z) = \frac{\left(\frac{L}{z}\right)^{3a} z \exp\left(-\frac{3z^2}{2R^2}\right) \left(2 \left(\frac{3z^2}{R^2}\right)^{\frac{3a-1}{4}} \mathbf{M}\left(\frac{-3a+1}{4}, \frac{3(-a+1)}{4}, \frac{3z^2}{R^2}\right) + 3(1-a)\exp\left(-\frac{3z^2}{2R^2}\right)\right)}{3(-1+3a)(-1+a)}$$

$$\mathbf{M}(\mu, \nu, z) = \exp\left(-\frac{z}{2}\right) z^{\frac{1}{2}+\nu} {}_1F_1\left(\frac{1}{2} + \nu - \mu, 1 + 2\nu, z\right)$$






Wich has maximal value at  $z_B \rightarrow \infty$  :  $S \rightarrow \frac{-F(z_A)}{2G_5}$

The entropy can be roughly estimate at  $a=1/2$  such as

$$S \sim E^{0.3} (1 + C_1 (\ln(E + 100))) - C_2$$

$$C_1 = -0.738, \quad C_2 = 0.393 \quad \text{at } 10 < E < 100 \quad \text{GeV}$$

$$C_1 = -0.073, \quad C_2 = 0.827 \quad \text{at } 100 < E < 1000 \quad \text{GeV}$$



# Conclusions

- The black holes formation in the domain wall-wall collisions is investigated in the wrapped  $AdS_5$  spaces.
- The several b-factor types: power-law, exponential and mixed are considered.
- The dependence of the entropy on the energy for different b-factors is analyzed.
- Our results (with the account of AdS/CFT-duality) allow to simulate the multiplicity dependence on the energy of the colliding heavy-ions in agreement with experimental data

$$b = (L/z)^a, \quad a \approx 0.47, \quad S \sim E^{0.3}, \quad S_{NN}^{0,15}$$

The additional logarithms appear when considering the mixed factor.



Thank you for attention!

