

The approach to gravity as a theory of embedded surface

Anton Sheykin
in collaboration with S.Paston

Saint-Petersburg State University

St. Petersburg 2013

Outline

- What is an isometrical embedding
- Applications
- Exact embeddings of the black holes
- The «extra solutions» of the Regge-Teitelboim equations
- Thermodynamical properties of the black holes

Idea of embedding

Janet-Cartan theorem (1916)

An arbitrary n -dimensional Riemannian manifold can be locally isometrically embedded in N -dimensional flat space with

$$N \geq \frac{n(n+1)}{2}. \quad (1)$$

For our 4D manifold $N = 10$; if the manifold has symmetries, N may be smaller. Metric of this manifold can be expressed in terms of embedding function:

$$g_{\mu\nu} = \partial_{\mu} y^a(x) \partial_{\nu} y^b(x) \eta_{ab}, \quad (2)$$

where $y^a(x)$ – embedding function, η_{ab} – metric of flat ambient space.

Idea of embedding

Janet-Cartan theorem (1916)

An arbitrary n -dimensional Riemannian manifold can be locally isometrically embedded in N -dimensional flat space with

$$N \geq \frac{n(n+1)}{2}. \quad (1)$$

For our 4D manifold $N = 10$; if the manifold has symmetries, N may be smaller. Metric of this manifold can be expressed in terms of embedding function:

$$g_{\mu\nu} = \partial_{\mu} y^a(x) \partial_{\nu} y^b(x) \eta_{ab}, \quad (2)$$

where $y^a(x)$ – embedding function, η_{ab} – metric of flat ambient space.

Idea of embedding

Janet-Cartan theorem (1916)

An arbitrary n -dimensional Riemannian manifold can be locally isometrically embedded in N -dimensional flat space with

$$N \geq \frac{n(n+1)}{2}. \quad (1)$$

For our 4D manifold $N = 10$; if the manifold has symmetries, N may be smaller. Metric of this manifold can be expressed in terms of embedding function:

$$g_{\mu\nu} = \partial_{\mu} y^a(x) \partial_{\nu} y^b(x) \eta_{ab}, \quad (2)$$

where $y^a(x)$ – embedding function, η_{ab} – metric of flat ambient space.

Embedding-based theory of gravity

Change of variables in action:

$$g_{\mu\nu}(x^\mu) \rightarrow y^a(x^\mu) \quad (3)$$

$$\begin{aligned} \delta S &= \frac{1}{2\kappa} \int d^4x \sqrt{-g} (G^{\mu\nu} - \kappa T^{\mu\nu}) \delta g_{\mu\nu} = \\ &= \frac{1}{2\kappa} \int d^4x \sqrt{-g} (G^{\mu\nu} - \kappa T^{\mu\nu}) \partial_\mu y^a(x) \partial_\nu \delta y^b(x) \eta_{ab} = \\ &= \partial_\mu (\sqrt{-g} (G^{\mu\nu} - \kappa T^{\mu\nu}) \partial_\nu y^a) = 0. \quad (4) \end{aligned}$$

(Regge, Teitelboim 1975) Natural appearance of the flat spacetime in this approach can be useful in the quantization of gravity:

- Preferred «time slicing»
- Definition of causality

Embedding-based theory of gravity

Change of variables in action:

$$g_{\mu\nu}(x^\mu) \rightarrow y^a(x^\mu) \quad (3)$$

$$\begin{aligned} \delta S &= \frac{1}{2\kappa} \int d^4x \sqrt{-g} (G^{\mu\nu} - \kappa T^{\mu\nu}) \delta g_{\mu\nu} = \\ &= \frac{1}{2\kappa} \int d^4x \sqrt{-g} (G^{\mu\nu} - \kappa T^{\mu\nu}) \partial_\mu y^a(x) \partial_\nu \delta y^b(x) \eta_{ab} = \\ &= \partial_\mu (\sqrt{-g} (G^{\mu\nu} - \kappa T^{\mu\nu}) \partial_\nu y^a) = 0. \quad (4) \end{aligned}$$

(Regge, Teitelboim 1975) Natural appearance of the flat spacetime in this approach can be useful in the quantization of gravity:

- Preferred «time slicing»
- Definition of causality

Embedding-based theory of gravity

Change of variables in action:

$$g_{\mu\nu}(x^\mu) \rightarrow y^a(x^\mu) \quad (3)$$

$$\begin{aligned} \delta S &= \frac{1}{2\kappa} \int d^4x \sqrt{-g} (G^{\mu\nu} - \kappa T^{\mu\nu}) \delta g_{\mu\nu} = \\ &= \frac{1}{2\kappa} \int d^4x \sqrt{-g} (G^{\mu\nu} - \kappa T^{\mu\nu}) \partial_\mu y^a(x) \partial_\nu \delta y^b(x) \eta_{ab} = \\ &= \partial_\mu (\sqrt{-g} (G^{\mu\nu} - \kappa T^{\mu\nu}) \partial_\nu y^a) = 0. \quad (4) \end{aligned}$$

(Regge, Teitelboim 1975) Natural appearance of the flat spacetime in this approach can be useful in the quantization of gravity:

- Preferred «time slicing»
- Definition of causality

Applications of isometrical embedding

Quantization

The Wheeler-de Witt equation for RT approach (Davidson, PRD 2003)

$$\frac{8\pi G}{2\sqrt{h}} \left[\left(\frac{\sqrt{h}}{8\pi G} \right)^2 (\hat{\lambda} + \mathcal{R}^{(3)})(x) - \right. \\ \left. - \hbar^2 \left((\Psi - \hat{\lambda}I)^{-1} \right)^{AB}(x) \frac{\delta^2}{\delta y^A(x) \delta y^B(x)} \right] \Phi[y] = 0. \quad (5)$$

Classification of exact solutions of Einstein equations

Embedding class $p = N - d = \frac{d(d-1)}{2}$ of a metric is invariant.

Geometrical properties of Riemannian manifolds

Fronsdal embedding of the Schwarzschild black hole is closely related to the Kruskal coordinates:

Applications of isometrical embedding

Quantization

The Wheeler-de Witt equation for RT approach (Davidson, PRD 2003)

$$\frac{8\pi G}{2\sqrt{h}} \left[\left(\frac{\sqrt{h}}{8\pi G} \right)^2 (\hat{\lambda} + \mathcal{R}^{(3)})(x) - \right. \\ \left. - \hbar^2 \left((\Psi - \hat{\lambda}I)^{-1} \right)^{AB} (x) \frac{\delta^2}{\delta y^A(x) \delta y^B(x)} \right] \Phi[y] = 0. \quad (5)$$

Classification of exact solutions of Einstein equations

Embedding class $p = N - d = \frac{d(d-1)}{2}$ of a metric is invariant.

Geometrical properties of Riemannian manifolds

Fronsdal embedding of the Schwarzschild black hole is closely related to the Kruskal coordinates:

Applications of isometrical embedding

Quantization

The Wheeler-de Witt equation for RT approach (Davidson, PRD 2003)

$$\frac{8\pi G}{2\sqrt{h}} \left[\left(\frac{\sqrt{h}}{8\pi G} \right)^2 (\hat{\lambda} + \mathcal{R}^{(3)})(x) - \right. \\ \left. - \hbar^2 \left((\Psi - \hat{\lambda}I)^{-1} \right)^{AB} (x) \frac{\delta^2}{\delta y^A(x) \delta y^B(x)} \right] \Phi[y] = 0. \quad (5)$$

Classification of exact solutions of Einstein equations

Embedding class $p = N - d = \frac{d(d-1)}{2}$ of a metric is invariant.


Geometrical properties of Riemannian manifolds

Fronsdal embedding of the Schwarzschild black hole is closely related to the Kruskal coordinates:

Fronsdal embedding (1959)

$$\begin{aligned} r > R : & & r < R : \\ y^0 &= 2R\sqrt{1 - \frac{R}{r}} \operatorname{sh}\left(\frac{t}{2R}\right), & y^0 &= \pm 2R\sqrt{\frac{R}{r} - 1} \operatorname{ch}\left(\frac{t}{2R}\right), \\ y^1 &= \pm 2R\sqrt{1 - \frac{R}{r}} \operatorname{ch}\left(\frac{t}{2R}\right), & y^1 &= 2R\sqrt{\frac{R}{r} - 1} \operatorname{sh}\left(\frac{t}{2R}\right), \\ & & y^2 &= r \cos(\theta), \\ & & y^3 &= r \sin(\theta) \cos(\phi), \\ & & y^4 &= r \sin(\theta) \sin(\phi), \\ & & y^5 &= g(r). \end{aligned} \tag{6}$$

Ambient space metric is $\eta_{ab} = \operatorname{diag}(1, -1, -1, -1, -1, -1)$.



Known embeddings of the Schwarzschild black hole

Kasner (1921)

$$y^0 = f(r), \quad y^1 = R\sqrt{1 - \frac{R}{r}} \sin(t/R), \quad y^2 = R\sqrt{1 - \frac{R}{r}} \cos(t/R). \quad (7)$$

Fujitani et al. (1961)

$$y^0 = t\sqrt{1 - \frac{R}{r}}, \quad y^{1,2} = \frac{1}{\sqrt{2}\gamma} \left(\frac{\gamma^2 t^2}{2} \mp 1 \right) \sqrt{1 - \frac{R}{r}} + \frac{u(r)}{\sqrt{2}}. \quad (8)$$

Davidson and Paz (1999)

$$y^{0,1} = \frac{R}{2\beta\sqrt{r_c r}} \left(e^{\beta t + u(r)} \mp \frac{r - r_c}{R} e^{-\beta t - u(r)} \right), \quad y^2 = kt. \quad (9)$$

$$y^3 = r \cos \theta, \quad y^4 = r \sin \theta \cos \phi, \quad y^5 = r \sin \theta \sin \phi.$$

New embeddings of the Schwarzschild black hole


Asymptotically flat embedding (S.Paston, A.S., CQG, 2012)

$$\begin{aligned}y^0 &= t', & y^1 &= \sqrt{\frac{27R^3}{r}} \sin \left(\frac{t'}{\sqrt{27R}} - \sqrt{\frac{(r+3R)^3}{27R^2r}} \right), \\y^2 &= \sqrt{\frac{27R^3}{r}} \cos \left(\frac{t'}{\sqrt{27R}} - \sqrt{\frac{(r+3R)^3}{27R^2r}} \right).\end{aligned}\quad (10)$$

Cubic embedding (S.P., A.S., TMsPh, 2013)

$$\begin{aligned}y^0 &= \frac{\xi^2}{6} t'^3 + \left(1 - \frac{R}{2r}\right) t' + u(r), \\y^1 &= \frac{\xi^2}{6} t'^3 - \frac{R}{2r} t' + u(r), \\y^2 &= \frac{\xi}{2} t'^2 + \frac{1}{2\xi} \left(1 - \frac{R}{r}\right).\end{aligned}\quad (11)$$

The global structure of the Schwarzschild black hole embeddings



I – our universe; II – black hole; III – white hole; IV – parallel universe.
Fronsdal – I,II,III,IV; Kasner, Fujitani – I; Davidson, As. flat, Cubic – I,II.

The «extra solutions» of the RT equations

$$\partial_{\mu}(\sqrt{-g}(G^{\mu\nu} - \kappa T^{\mu\nu})\partial_{\nu}y^a) = 0. \quad (12)$$

It looks like conservation law for some current:

$$\partial_{\mu}j^{a\mu} = 0; \quad j^{a\mu} = \sqrt{-g}(G^{\mu\nu} - \kappa T^{\mu\nu})\partial_{\nu}y^a. \quad (13)$$

RT equations are equivalent to Einstein equations with additional «matter» $\tau^{\mu\nu}$

$$G_{\mu\nu} - \kappa T_{\mu\nu} = \frac{j_{a\nu}\partial_{\mu}y^a}{\sqrt{-g}} \equiv \kappa\tau_{\mu\nu}, \quad (14)$$

which dynamics is governed by equation

$$\partial_{\mu}(\sqrt{-g}\tau^{\mu\nu}\partial_{\nu}y^a) = 0. \quad (15)$$

The «extra solutions» of the RT equations

$$\partial_\mu(\sqrt{-g}(G^{\mu\nu} - \kappa T^{\mu\nu})\partial_\nu y^a) = 0. \quad (12)$$

It looks like conservation law for some current:

$$\partial_\mu j^{a\mu} = 0; \quad j^{a\mu} = \sqrt{-g}(G^{\mu\nu} - \kappa T^{\mu\nu})\partial_\nu y^a. \quad (13)$$

RT equations are equivalent to Einstein equations with additional «matter» $\tau^{\mu\nu}$

$$G_{\mu\nu} - \kappa T_{\mu\nu} = \frac{j_{a\nu}\partial_\mu y^a}{\sqrt{-g}} \equiv \kappa\tau_{\mu\nu}, \quad (14)$$

which dynamics is governed by equation

$$\partial_\mu(\sqrt{-g}\tau^{\mu\nu}\partial_\nu y^a) = 0. \quad (15)$$

The «extra solutions» of the RT equations

$$\partial_{\mu}(\sqrt{-g}(G^{\mu\nu} - \kappa T^{\mu\nu})\partial_{\nu}y^a) = 0. \quad (12)$$

It looks like conservation law for some current:

$$\partial_{\mu}j^{a\mu} = 0; \quad j^{a\mu} = \sqrt{-g}(G^{\mu\nu} - \kappa T^{\mu\nu})\partial_{\nu}y^a. \quad (13)$$

RT equations are equivalent to Einstein equations with additional «matter» $\tau^{\mu\nu}$

$$G_{\mu\nu} - \kappa T_{\mu\nu} = \frac{j_{a\nu}\partial_{\mu}y^a}{\sqrt{-g}} \equiv \kappa\tau_{\mu\nu}, \quad (14)$$

which dynamics is governed by equation

$$\partial_{\mu}(\sqrt{-g}\tau^{\mu\nu}\partial_{\nu}y^a) = 0. \quad (15)$$

$$\partial_\mu(\sqrt{-g}\tau^{\mu\nu}\partial_\nu y^a) = 0. \quad (16)$$

If $\tau^{\mu\nu} = 0$ in some boundary or Cauchy surface, RT equations become equivalent Einstein equations permanently (Paston, Franke, 2007). In some cases $\tau^{\mu\nu}$ vanishes automatically:

- Inflation in early universe (in assumption of Friedmann symmetry existence):

$j^{a\nu}$ is conserved, \sqrt{g} increases exponentially $\Rightarrow \tau^{\mu\nu}$ decreases exponentially (Paston, A.S., 2012).

- Vacuum spherically symmetric static case:

In all 6 types of embedding exists y^* such that $\partial_\nu y^*$ depend only on r .

$$\begin{aligned} \partial_\mu(\partial_\nu y^* \sqrt{-g} G^{\mu\nu}) &= \partial_r(\partial_r y^* G^{rr} \sqrt{-g}) \Rightarrow \\ &\Rightarrow \partial_r y^* G^{rr} \sqrt{-g} = C. \end{aligned} \quad (17)$$

It can be shown that $\partial_r y^* G^{rr} \sqrt{-g} \rightarrow 0$ when $r \rightarrow \infty$

$$\Rightarrow C = 0 \Rightarrow G^{rr} = 0 \Rightarrow G^{\mu\nu} = 0.$$

$$\partial_\mu(\sqrt{-g}\tau^{\mu\nu}\partial_\nu y^a) = 0. \quad (16)$$

If $\tau^{\mu\nu} = 0$ in some boundary or Cauchy surface, RT equations become equivalent Einstein equations permanently (Paston, Franke, 2007). In some cases $\tau^{\mu\nu}$ vanishes automatically:

- Inflation in early universe (in assumption of Friedmann symmetry existence):

$j^{a\nu}$ is conserved, \sqrt{g} increases exponentially $\Rightarrow \tau^{\mu\nu}$ decreases exponentially (Paston, A.S., 2012).

- Vacuum spherically symmetric static case:

In all 6 types of embedding exists y^* such that $\partial_\nu y^*$ depend only on r .

$$\begin{aligned} \partial_\mu(\partial_\nu y^* \sqrt{-g} G^{\mu\nu}) &= \partial_r(\partial_r y^* G^{rr} \sqrt{-g}) \Rightarrow \\ &\Rightarrow \partial_r y^* G^{rr} \sqrt{-g} = C. \end{aligned} \quad (17)$$

It can be shown that $\partial_r y^* G^{rr} \sqrt{-g} \rightarrow 0$ when $r \rightarrow \infty$
 $\Rightarrow C = 0 \Rightarrow G^{rr} = 0 \Rightarrow G^{\mu\nu} = 0.$

$$\partial_\mu(\sqrt{-g}\tau^{\mu\nu}\partial_\nu y^a) = 0. \quad (16)$$

If $\tau^{\mu\nu} = 0$ in some boundary or Cauchy surface, RT equations become equivalent Einstein equations permanently (Paston, Franke, 2007). In some cases $\tau^{\mu\nu}$ vanishes automatically:

- Inflation in early universe (in assumption of Friedmann symmetry existence):

$j^{a\nu}$ is conserved, \sqrt{g} increases exponentially $\Rightarrow \tau^{\mu\nu}$ decreases exponentially (Paston, A.S., 2012).

- Vacuum spherically symmetric static case:

In all 6 types of embedding exists y^* such that $\partial_\nu y^*$ depend only on r .

$$\begin{aligned} \partial_\mu(\partial_\nu y^* \sqrt{-g} G^{\mu\nu}) &= \partial_r(\partial_r y^* G^{rr} \sqrt{-g}) \Rightarrow \\ &\Rightarrow \partial_r y^* G^{rr} \sqrt{-g} = C. \end{aligned} \quad (17)$$

It can be shown that $\partial_r y^* G^{rr} \sqrt{-g} \rightarrow 0$ when $r \rightarrow \infty$
 $\Rightarrow C = 0 \Rightarrow G^{rr} = 0 \Rightarrow G^{\mu\nu} = 0.$

$$\partial_\mu(\sqrt{-g}\tau^{\mu\nu}\partial_\nu y^a) = 0. \quad (16)$$

If $\tau^{\mu\nu} = 0$ in some boundary or Cauchy surface, RT equations become equivalent Einstein equations permanently (Paston, Franke, 2007). In some cases $\tau^{\mu\nu}$ vanishes automatically:

- Inflation in early universe (in assumption of Friedmann symmetry existence):

$j^{a\nu}$ is conserved, \sqrt{g} increases exponentially $\Rightarrow \tau^{\mu\nu}$ decreases exponentially (Paston, A.S., 2012).

- Vacuum spherically symmetric static case:

In all 6 types of embedding exists y^* such that $\partial_\nu y^*$ depend only on r .

$$\begin{aligned} \partial_\mu(\partial_\nu y^* \sqrt{-g} G^{\mu\nu}) &= \partial_r(\partial_r y^* G^{rr} \sqrt{-g}) \Rightarrow \\ &\Rightarrow \partial_r y^* G^{rr} \sqrt{-g} = C. \end{aligned} \quad (17)$$

It can be shown that $\partial_r y^* G^{rr} \sqrt{-g} \rightarrow 0$ when $r \rightarrow \infty$
 $\Rightarrow C = 0 \Rightarrow G^{rr} = 0 \Rightarrow G^{\mu\nu} = 0.$

New global embeddings of the charged black hole

$$ds^2 = \left(1 - \frac{2m}{r} + \frac{q^2}{r^2}\right) dt^2 - \frac{dr^2}{\left(1 - \frac{2m}{r} + \frac{q^2}{r^2}\right)} - r^2 d\Omega^2.$$

Spiral

$$\begin{aligned}y^0 &= \frac{\sqrt{(mr - q^2)^2 + b^2 r^2}}{\alpha q r} \sin(\alpha t' + u(r)), \\y^1 &= \frac{\sqrt{(mr - q^2)^2 + b^2 r^2}}{\alpha q r} \cos(\alpha t' + u(r)), \\y^2 &= \frac{\sqrt{b^2 + m^2 - q^2}}{q} t'.\end{aligned}\tag{18}$$

Exponential


$$y^0 = \gamma t',$$
$$y^{1,2} = \frac{e^{-\beta t' - v(r)}}{2\beta} \mp \left(1 - \frac{2m}{r} + \frac{q^2}{r^2} - \gamma^2\right) \frac{e^{\beta t' + v(r)}}{2\beta}.$$

Cubic

$$y^0 = -\frac{q^4}{8m^3} \left(1 - \frac{2m}{r} + \frac{q^2}{r^2}\right) + \frac{2m^3}{q^4} t'^2,$$
$$y^{1,2} = w(r) - \frac{1}{4} \left(1 - \frac{2m}{r} + \frac{q^2}{r^2}\right) t' + \frac{4m^6}{3q^8} t'^3 \mp t'.$$

Ambient space metric is $\eta_{ab} = \text{diag}(1, 1, -1, -1, -1, -1)$.

The global structure of the charged black hole embeddings



New global embeddings of the black hole in de Sitter space

$$ds^2 = \left(1 - \frac{2m}{r} - \frac{\Lambda r^2}{3}\right) dt^2 - \frac{dr^2}{\left(1 - \frac{2m}{r} - \frac{\Lambda r^2}{3}\right)} - r^2 d\Omega^2.$$

Cubic

$$y^{0,1} = \frac{\xi^2}{6} t'^3 + \frac{g_{00} \pm 1}{2} t' + u(r), \quad y^2 = \frac{\xi}{2} t'^2 + \frac{g_{00}}{2\xi}. \quad (19)$$

Spiral

$$y^0 = t', \quad y^1 = \sqrt{\frac{\Lambda r^3 + 2m}{\alpha^2 r}} \sin(\alpha t' + w(r)), \quad y^2 = \sqrt{\frac{\Lambda r^3 + 2m}{\alpha^2 r}} \cos(\alpha t' + w(r)). \quad (20)$$

Ambient space metric is $\eta_{ab} = \text{diag}(1, -1, -1, -1, -1, -1)$.

Hawking and Unruh effect

Hawking effect

A black hole has a radiation with a thermal spectrum $T_0 = \frac{1}{4\pi R}$.

Tolman law

In thermal equilibrium a temperature is not constant when space is curved. For the Schwarzschild black hole

$$T = T_0 / \sqrt{1 - R/r}. \quad (21)$$

Unruh effect

Uniformly accelerated observer, when coupled to the quantum fields, detects a radiation with a thermal spectrum and a temperature proportional to his acceleration: $T = \frac{a}{2\pi}$.

Hawking and Unruh effect

Hawking effect

A black hole has a radiation with a thermal spectrum $T_0 = \frac{1}{4\pi R}$.

Tolman law

In thermal equilibrium a temperature is not constant when space is curved. For the Schwarzschild black hole

$$T = T_0 / \sqrt{1 - R/r}. \quad (21)$$

Unruh effect

Uniformly accelerated observer, when coupled to the quantum fields, detects a radiation with a thermal spectrum and a temperature proportional to his acceleration: $T = \frac{a}{2\pi}$.

Hawking and Unruh effect

Hawking effect

A black hole has a radiation with a thermal spectrum $T_0 = \frac{1}{4\pi R}$.

Tolman law

In thermal equilibrium a temperature is not constant when space is curved. For the Schwarzschild black hole

$$T = T_0 / \sqrt{1 - R/r}. \quad (21)$$

Unruh effect

Uniformly accelerated observer, when coupled to the quantum fields, detects a radiation with a thermal spectrum and a temperature proportional to his acceleration: $T = \frac{a}{2\pi}$.

Connection between Hawking and Unruh effect in ambient space


Deser & Levin, PRD 1999

Unruh radiation detected by observer moving on the embedding surface in the ambient space has the same spectrum and temperature as the Hawking radiation from the horizon in the corresponding manifold: $T = \frac{a_{emb}}{2\pi}$. This hypothesis was tested for Fronsdal embedding.

Further development

- Hong, GRG 2003 – charged black holes,
- Chen et al., JHEP 2004 – stationary motions,
- Santos et al., PRD 2004 – D -dimensional black holes,
- ...

$$T = a/2\pi = \frac{1}{4\pi R\sqrt{1-R/r}}, \quad T_0 = T\sqrt{1-R/r} = \text{const.} \quad (22)$$



Accelerations for new embeddings

- Asymptotically flat: $a = \frac{1}{\sqrt{27Rr} \left(1 - \frac{R}{r}\right)}$
- Davidson-Paz: $a = \frac{\beta}{1 - \frac{R}{r}} \sqrt{1 - \frac{R}{r} + k^2}$
- Cubic $a = \frac{\xi}{\left(1 - \frac{R}{r}\right)}$

Corresponding temperatures (if we assume that the temperature and acceleration are always related as in the Unruh effect) violate the Tolman law and do not coincide with BH temperature!

For moving on the arbitrary trajectories the spectrum of Unruh radiation is not exactly thermal, but at sufficiently slowly varying accelerations ($\dot{a}/a^2 \ll 1$) the Unruh formula works good (Barbado & Visser, PRD, 2012).

Exact spectrum was found to be non-thermal for the trajectories corresponding to lines of time in cubic (Letaw, PRD 1981) and Davidson-Paz (Abdolrahimi, arXiv:1304:4237) embeddings.

For moving on the arbitrary trajectories the spectrum of Unruh radiation is not exactly thermal, but at sufficiently slowly varying accelerations ($\dot{a}/a^2 \ll 1$) the Unruh formula works good (Barbado & Visser, PRD, 2012).

Exact spectrum was found to be non-thermal for the trajectories corresponding to lines of time in cubic (Letaw, PRD 1981) and Davidson-Paz (Abdolrahimi, arXiv:1304:4237) embeddings.

Summary

- Isometrical embedding is a powerful tool for studying of Riemannian manifolds.
- Some features of embeddings possibly can help to quantize gravity.
- In some physically interesting cases the predictions of the embedding-based theory of gravity coincide with the predictions of the Einstein theory.
- Exact embeddings of black holes might be related to their thermodynamical properties.
- Mapping between Hawking effect and Unruh effect in ambient space holds only for hyperbolic (Fronsdal-like) embeddings.

Summary

- Isometrical embedding is a powerful tool for studying of Riemannian manifolds.
- Some features of embeddings possibly can help to quantize gravity.
- In some physically interesting cases the predictions of the embedding-based theory of gravity coincide with the predictions of the Einstein theory.
- Exact embeddings of black holes might be related to their thermodynamical properties.
- Mapping between Hawking effect and Unruh effect in ambient space holds only for hyperbolic (Fronsdal-like) embeddings.

Summary

- Isometrical embedding is a powerful tool for studying of Riemannian manifolds.
- Some features of embeddings possibly can help to quantize gravity.
- In some physically interesting cases the predictions of the embedding-based theory of gravity coincide with the predictions of the Einstein theory.
- Exact embeddings of black holes might be related to their thermodynamical properties.
- Mapping between Hawking effect and Unruh effect in ambient space holds only for hyperbolic (Fronsdal-like) embeddings.

Summary

- Isometrical embedding is a powerful tool for studying of Riemannian manifolds.
- Some features of embeddings possibly can help to quantize gravity.
- In some physically interesting cases the predictions of the embedding-based theory of gravity coincide with the predictions of the Einstein theory.
- Exact embeddings of black holes might be related to their thermodynamical properties.
- Mapping between Hawking effect and Unruh effect in ambient space holds only for hyperbolic (Fronsdal-like) embeddings.

Summary

- Isometrical embedding is a powerful tool for studying of Riemannian manifolds.
- Some features of embeddings possibly can help to quantize gravity.
- In some physically interesting cases the predictions of the embedding-based theory of gravity coincide with the predictions of the Einstein theory.
- Exact embeddings of black holes might be related to their thermodynamical properties.
- Mapping between Hawking effect and Unruh effect in ambient space holds only for hyperbolic (Fronsdal-like) embeddings.